

BOUND FOR THE NUMBER OF 1-DIMENSIONAL FIBERS OF A PROJECTIVE MORPHISM

TRAN QUANG HOA

ABSTRACT. Given a birational parameterization $\phi : \mathbb{P}_k^2 \dashrightarrow \mathbb{P}_k^3$ of an algebraic surface $\mathcal{S} \subset \mathbb{P}_k^3$, we bound the number of 1-dimensional fibers of the canonical projection of the graph of ϕ onto its image.

Keyword: *Fibers of morphism, implicitization, implicit equation, parameterization surfaces, elimination theory, approximation complex, geometric modeling.*

1. INTRODUCTION

Rational algebraic surfaces can be described in several ways, the most common being parametric and implicit representations. Parametric representations describe such surface as the image of a rational map, whereas implicit representations describe it as the zero locus of a certain algebraic equation, e.g. as the zeros of a polynomial. Both representations have a wide range of applications in Computer Aided Geometric Design (CAGD), and depending on the problem one needs to solve, one or the other might be better suited. It is thus interesting to be able to pass from parametric representations to implicit equations. This is a classical problem and there are numerous approaches to its solution. For a good historical overview on this subject we refer the reader to [SC95] and [Cox01].

The implicitization problem has been widely studied and is always an active research area. It is basically an elimination problem, and thus can be solved by methods based on Gröbner basis computations. However, it is known to be quite slow in practice and hence is rarely used in geometric modeling (see, e.g., [Hof89]). A more common method for finding the implicit equation is to eliminate by computing the resultant of the polynomials (see, e.g., [Jou96, Section 5.3.17], [Bus01]). But in many applications, the resultant vanishes identically due to the presence of base points. In consequence, the search of formulas for implicitization rational surfaces with base points is a very active area of research due to the fact that, in practical industrial design, base points show up quite frequently. In [MC92], a perturbation is applied to resultants in order to obtain a nonzero multiple of the implicit equation. Many other approaches have been done in this direction. Lately, it is shown how to compute the implicit equation as the determinant of the approximation complexes (see, e.g., [BJ03], [BCD03], [BC05], [Cha06], [BCJ09], ...).

In geometric modeling, it is of interest to determine the singularities of the parametrization, that is, to know whether a point is being painted twice, or not.

It is important to give a computationally efficient criterion to answer this question. The matrix representations are also useful for solving this problem (see, e.g., [BBC14], [Bus14]).

The main goal of this paper is to estimate the number of the particular points on a surface. More precisely, let k a field and let ϕ be a parameterization from $\mathbb{P}_k^2$ to $\mathbb{P}_k^3$. Let $\Gamma \subset \mathbb{P}_k^2 \times \mathbb{P}_k^3$ be the closure of the graph of ϕ and $\pi_2 : \mathbb{P}_k^2 \times \mathbb{P}_k^3 \longrightarrow \mathbb{P}_k^3$ the projection. For every $\mathbf{p} \in \mathbb{P}_k^3$, the fiber of π_2 at $\mathbf{p}$ is the subscheme

$$\pi_2^{-1}(\mathbf{p}) := \Gamma \times_{\mathbb{P}_k^3} \text{Spec}(k(\mathbf{p})) \subset \mathbb{P}_{k(\mathbf{p})}^2$$

where $k(\mathbf{p})$ its residue field. We denote by

$$\mathcal{Y} := \{\mathbf{p} \in \mathbb{P}_k^3 \mid \dim \pi_2^{-1}(\mathbf{p}) = 1\}.$$

In this paper we will propose a method to estimate the cardinality of $\mathcal{Y}$. For any $\mathbf{p} \in \mathcal{Y}$, we denote by $h_{\mathbf{p}} \in R$ the defining equation of the unmixed component of the fiber $\pi_2^{-1}(\mathbf{p})$.

The main theorem is the following.

Theorem. *Let $I = (f_0, \dots, f_3)$ be a homogeneous ideal of $R := k[X_1, X_2, X_3]$ generated by four homogeneous polynomials of the same degree d , not all zero. Let $I^{\text{sat}} := I :_R (X_1, X_2, X_3)^\infty$ be the saturation of I . Suppose that $\inf\{\nu \mid I_\nu^{\text{sat}} \neq 0\} = d$ and $\mathcal{P} = \text{Proj}(R/I)$ is locally a complete intersection of dimension zero.*

(i) *If $d = 3$, then*

$$\sum_{\mathbf{p} \in \mathcal{Y}} \deg(h_{\mathbf{p}}) \leq 4.$$

(ii) *If $d \geq 4$, then*

$$\sum_{\mathbf{p} \in \mathcal{Y}} \deg(h_{\mathbf{p}}) \leq \left\lfloor \frac{d}{2} \right\rfloor d - 1.$$

It was noticed in [BBC14] that if $I_\nu^{\text{sat}} \neq 0$ for some $\nu < d$ then $\sum_{\mathbf{p} \in \mathcal{Y}} \deg(h_{\mathbf{p}}) \leq \nu$.

2. PRELIMINARIES

Let $R := k[X_1, X_2, X_3]$ be a polynomial ring over the field k with its standard $\mathbb{Z}$ -grading $\deg(X_i) = 1$ for all $i = 1, 2, 3$. Suppose we are given an integer $d \geq 1$ and four homogeneous polynomials $f_0, \dots, f_3 \in R_d$, not all zero. We denote by I the ideal of R generated by these polynomials and set $\mathcal{P} := \text{Proj}(R/I) \subseteq \mathbb{P}_k^2 := \text{Proj}(R)$. Let $B := k[T_0, T_1, T_2, T_3]$ with its standard grading and consider the rational map

$$\begin{aligned} \phi : \mathbb{P}_k^2 &\dashrightarrow \mathbb{P}_k^3 \\ x &\longmapsto (f_0(x) : \dots : f_3(x)). \end{aligned}$$

We will always assume throughout this paper that ϕ is generically finite onto its image, or equivalently that the closed image of ϕ is a surface $\mathcal{S}$ in $\mathbb{P}_k^3 := \text{Proj}(B)$, and that the base locus $\mathcal{P}$ of ϕ is supported on a finite set of points.

Let $\Gamma_0 \subset \mathbb{P}_k^2 \times \mathbb{P}_k^3$ be the graph of $\phi : \mathbb{P}_k^2 \setminus \mathcal{P} \longrightarrow \mathbb{P}_k^3$ and Γ the Zariski closure of Γ_0 . We have the following diagram

$$\begin{array}{ccc} \Gamma & \xrightarrow{\quad} & \mathbb{P}_k^2 \times \mathbb{P}_k^3 \\ \pi_1 \downarrow & \searrow \pi_2 & \\ \mathbb{P}_k^2 & \xrightarrow[\phi]{} & \mathbb{P}_k^3 \end{array}$$

where the maps π_1, π_2 are the canonical projections. One has

$$\mathcal{S} = \overline{\pi_2(\Gamma_0)} = \pi_2(\Gamma)$$

where the bar denotes the Zariski closure.

Furthermore Γ is the irreducible subscheme of $\mathbb{P}_k^2 \times \mathbb{P}_k^3$ defined by the Rees algebra $\mathcal{R}_{\mathcal{I}} := \text{Rees}_R(I)$ (see [Har77, Chapter II, §7]). Let $S := R \otimes_k B = R[T_0, \dots, T_3]$ with the standard bigraded structure by the canonical grading $\deg(X_i) = (1, 0)$ and $\deg(T_j) = (0, 1)$ for all $i = 1, 2, 3$ and $j = 0, \dots, 3$. The natural bigraded morphism of bigraded k -algebras

$$\begin{aligned} \alpha : S &\longrightarrow \mathcal{R}_{\mathcal{I}} = \bigoplus_{s \geq 0} I(d)^s = \bigoplus_{s \geq 0} I^s(sd) \\ T_i &\longmapsto f_i \end{aligned}$$

is onto and corresponds to the embedding $\Gamma \subset \mathbb{P}_k^2 \times \mathbb{P}_k^3$.

Let $\mathfrak{P}$ be the kernel of α . Then it is a homogeneous ideal of S and the part of degree one of $\mathfrak{P}$ in T_i , denoted by $\mathfrak{P}_1 = \mathfrak{P}_{(*,1)}$, is the module of syzygies of the f_i

$$a_0 T_0 + \dots + a_3 T_3 \in \mathfrak{P}_1 \iff a_0 f_0 + \dots + a_3 f_3 = 0.$$

Set $\mathcal{S}_{\mathcal{I}} := \text{Sym}_R(I)$ for the symmetric algebra of I . The natural bigraded epimorphisms

$$S \longrightarrow S/(\mathfrak{P}_1) \simeq \mathcal{S}_{\mathcal{I}} \quad \text{and} \quad \mathcal{S}_{\mathcal{I}} \simeq S/(\mathfrak{P}_1) \longrightarrow S/\mathfrak{P} \simeq \mathcal{R}_{\mathcal{I}}$$

correspond to the embeddings of schemes $\Gamma \subseteq V \subset \mathbb{P}_k^2 \times \mathbb{P}_k^3$ where V is the projective scheme defined by $\mathcal{S}_{\mathcal{I}}$.

Let K be defined by the following exact sequence

$$0 \longrightarrow K \longrightarrow \mathcal{S}_{\mathcal{I}} \longrightarrow \mathcal{R}_{\mathcal{I}} \longrightarrow 0.$$

Notice that the module K is supported in $\mathcal{P}$ because I is locally trivial outside $\mathcal{P}$.

Let us now denote by I^{sat} the ideal

$$I :_R \mathfrak{m}^\infty = \bigcup_{n \in \mathbb{N}} I :_R \mathfrak{m}^n$$

where $\mathfrak{m} := (X_1, X_2, X_3)$ is the irrelevant maximal ideal of R .

Finally, we recall that for a nonzero $\mathbb{Z}$ -graded R -module of finite type M , its initial degree is defined by

$$\text{indeg}(M) := \inf\{\nu \mid M_\nu \neq 0\},$$

with the convention $\text{indeg}(M) = +\infty$ when $M = 0$.

3. FIBERS OF THE CANONICAL PROJECTION OF THE GRAPH OF ϕ ONTO $\mathcal{S}$

From now on we assume that the ideal I is locally a complete intersection outside $V(\mathfrak{m})$. Under this hypothesis, the symmetric algebra $\mathcal{S}_{\mathcal{I}}$ is projectively isomorphic to the Rees algebra $\mathcal{R}_{\mathcal{I}}$ (see for instance [BC05, Theorem 4]). In other words, the graph $\Gamma \subset \mathbb{P}_k^2 \times \mathbb{P}_k^3$ of the parameterization ϕ is equal to the subscheme V of $\mathbb{P}_k^2 \times \mathbb{P}_k^3$ defined by $\mathcal{S}_{\mathcal{I}}$. For every $\mathfrak{p} \in \mathbb{P}_k^3$, we will denote by $k(\mathfrak{p})$ its residue field $B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$. The fiber of π_2 at $\mathfrak{p}$ is the subscheme

$$\pi_2^{-1}(\mathfrak{p}) = \text{Proj}(\mathcal{R}_{\mathcal{I}} \otimes_B k(\mathfrak{p})) = \text{Proj}(\mathcal{S}_{\mathcal{I}} \otimes_B k(\mathfrak{p})) \subset \mathbb{P}_{k(\mathfrak{p})}^2.$$

In [BBC14], it is shown that π_2 only admits fibers of dimension 0 or 1. Furthermore, it is easily seen that the number of 1-dimensional fibers is finite. Our purpose is to bound the cardinality of the set of points in $\mathbb{P}_k^3$ with a 1-dimensional fiber, that is, the cardinality of the set

$$\mathcal{Y} = \{\mathfrak{p} \in \mathbb{P}_k^3 \mid \dim \pi_2^{-1}(\mathfrak{p}) = 1\}.$$

For any $\mathfrak{p} \in \mathcal{Y}$, we will denote by $h_{\mathfrak{p}} \in R$ the defining equation of the unmixed component of the fiber $\pi_2^{-1}(\mathfrak{p})$. We begin with an important remark in this direction.

Proposition 1. *If there exists an integer s such that $\nu = \text{indeg}((I^s)^{\text{sat}}) < sd$, then*

$$\sum_{\mathfrak{p} \in \mathcal{Y}} \deg(h_{\mathfrak{p}}) \leq \nu < sd.$$

Proof. First notice that $\mathcal{Y}$ is finite. By [BBC14, Lemma 10], there exists a homogeneous polynomial $f \in I$ of degree d such that for any $\mathfrak{p} \in \mathcal{Y}$, we have

$$I = (f) + h_{\mathfrak{p}}(g_{1\mathfrak{p}}, g_{2\mathfrak{p}}, g_{3\mathfrak{p}}) \quad \text{and} \quad I^{\text{sat}} \subseteq (f, h_{\mathfrak{p}})$$

for some $g_{1\mathfrak{p}}, g_{2\mathfrak{p}}, g_{3\mathfrak{p}} \in R$.

Since $(f, h_{\mathfrak{p}})$ is a complete intersection ideal, it follows from [ZS60, Lemma 5, Appendix 6] that $(f, h_{\mathfrak{p}})^s$ is unmixed, hence saturated. Therefore for all $\mathfrak{p} \in \mathcal{Y}$, we have

$$(I^s)^{\text{sat}} \subseteq ((I^{\text{sat}})^s)^{\text{sat}} \subseteq ((f, h_{\mathfrak{p}})^s)^{\text{sat}} = (f, h_{\mathfrak{p}})^s = (f^s, f^{s-1}h_{\mathfrak{p}}, \dots, h_{\mathfrak{p}}^s).$$

If $F \in (I^s)^{\text{sat}}$ such that $\deg(F) = \nu < sd$, then $h_{\mathfrak{p}}$ is a divisor of F . Moreover, if $\mathfrak{p} \neq \mathfrak{p}'$ in $\mathcal{Y}$, then $\gcd(h_{\mathfrak{p}}, h_{\mathfrak{p}'}) = 1$. We deduce that

$$\prod_{\mathfrak{p} \in \mathcal{Y}} h_{\mathfrak{p}} \mid F$$

which gives

$$\sum_{\mathfrak{p} \in \mathcal{Y}} \deg(h_{\mathfrak{p}}) \leq \deg(F) = \nu < sd.$$

□

In particulier if $\text{indeg}(I^{\text{sat}}) = \delta < d$, by Proposition 1, then

$$\sum_{\mathfrak{p} \in \mathcal{Y}} \deg(h_{\mathfrak{p}}) \leq \delta < d.$$

From now on we want to bound the cardinality of the set $\mathcal{Y}$, the delicate case is when the ideal I satisfies $\text{indeg}(I^{\text{sat}}) = \text{indeg}(I) = d$.

Example 2. Consider the parameterization given by

$$\begin{aligned} f_0 &= X_2^2 X_3^4 - X_2^4 X_3^2, \\ f_1 &= X_1^4 X_3^2 - X_3^6, \\ f_2 &= X_1^2 X_2^2 X_3^2 - X_1^2 X_2^4, \\ f_3 &= X_1^4 X_2^2 - X_2^2 X_3^4. \end{aligned}$$

Using Macaulay2 [Mac2], we see that I is a saturated ideal with $\mathcal{P}$ locally a complete intersection of dimension zero. Since $\text{indeg}((I^2)^{\text{sat}}) = 8 < 2.6 = 12$, Proposition 1 shows that

$$\sum_{\mathfrak{p} \in \mathcal{Y}} \deg(h_{\mathfrak{p}}) \leq 8.$$

Precisely, I admits a minimal free resolution of the form

$$0 \longrightarrow R(-8)^3 \xrightarrow{M} R(-6)^4 \longrightarrow R \longrightarrow R/I \longrightarrow 0$$

where the matrix M is given by

$$\begin{pmatrix} -X_2^2 + X_3^2 & -X_3^2 & 0 \\ X_1^2 & 0 & -X_3^2 \\ 0 & X_2^2 & 0 \\ -X_3^2 & 0 & X_1^2 \end{pmatrix}.$$

We thus get $\mathcal{Y} = \{\mathfrak{p}_1, \mathfrak{p}_2, \mathfrak{p}_3, \mathfrak{p}_4\}$ with

$$\begin{aligned} \mathfrak{p}_1 &= (1 : 0 : 1 : 0), & h_{\mathfrak{p}_1} &= X_2^2 - X_3^2, & \mathfrak{p}_2 &= (0 : 1 : 0 : 1), & h_{\mathfrak{p}_2} &= X_1^2 - X_3^2, \\ \mathfrak{p}_3 &= (0 : 1 : 0 : -1), & h_{\mathfrak{p}_3} &= X_1^2 + X_3^2, & \mathfrak{p}_4 &= (0 : 0 : 1 : 0), & h_{\mathfrak{p}_4} &= X_2^2, \end{aligned}$$

which implies

$$\sum_{\mathfrak{p} \in \mathcal{Y}} \deg(h_{\mathfrak{p}}) = 8.$$

4. ESTIMATION OF NUMBER OF 1-DIMENSIONAL FIBERS OF THE CANONICAL PROJECTION OF THE GRAPH OF ϕ ONTO $\mathcal{S}$

As we already noticed that $\sum_{\mathfrak{p} \in \mathcal{Y}} \deg(h_{\mathfrak{p}}) < d$ if $\text{indeg}(I^{\text{sat}}) < d$, we assume that $\text{indeg}(I^{\text{sat}}) = \text{indeg}(I) = d$.

Recall that $\mathcal{P} = \text{Proj}(R/I)$ is assumed to be locally a complete intersection of dimension zero. Under this hypothesis, the module K is supported in $\mathfrak{m}S$, hence $H_{\mathfrak{m}}^i(K) = 0$ for any $i \geq 1$. We deduce that

$$H_{\mathfrak{m}}^i(\mathcal{S}_{\mathcal{I}}) \simeq H_{\mathfrak{m}}^i(\mathcal{R}_{\mathcal{I}}), \forall i \geq 1.$$

Here we denote the i -th local cohomology modules of a R -module M with support in the irrelevant maximal ideal $\mathfrak{m} = (X_1, X_2, X_3)$ by $H_{\mathfrak{m}}^i(M)$. We set

$$a_i(M) := \sup\{\mu \mid H_{\mathfrak{m}}^i(M)_{\mu} \neq 0\},$$

with the convention $a_i(M) := -\infty$ when $H_{\mathfrak{m}}^i(M) = 0$. Recall that the Castelnuovo-Mumford regularity of M is defined by

$$\text{reg}(M) := \max_i \{a_i(M) + i\}.$$

Let $K_{\bullet} := K_{\bullet}(\mathbf{f}; R)$ be the Koszul complex associated to the sequence of polynomials $\mathbf{f} := (f_0, f_1, f_2, f_3)$ in R . We denote by $Z_i := Z_i(\mathbf{f}; R)$, $B_i := B_i(\mathbf{f}; R)$, $H_i :=$

$H_i(\mathbf{f}; R) = Z_i/B_i$ the module of cycles, the module of boundaries and the homology module in degree i , respectively (see [BJ03, §4] or [BC05, §3] for an introduction to these complexes in this context). Since the ideal I is homogeneous, these modules inherit a natural structure of graded R -modules. Let $\mathcal{Z}_\bullet$ be the approximation complex of cycles associated to the ideal $I = (\mathbf{f})$. By definition $\mathcal{Z}_q = Z_q[qd] \otimes_R R[T_0, \dots, T_3](-q)$ for all $q = 0, \dots, 3$. This complex is of the form

$$(\mathcal{Z}_\bullet) : \quad 0 \longrightarrow \mathcal{Z}_3 \xrightarrow{v_3} \mathcal{Z}_2 \xrightarrow{v_2} \mathcal{Z}_1 \xrightarrow{v_1} \mathcal{Z}_0 = R[T_0, \dots, T_3] \xrightarrow{v_0} 0$$

where $v_1(a_0, a_1, a_2, a_3) = a_0T_0 + \dots + a_3T_3$. It is proved in [BC05, Theorem 4] that the complex $(\mathcal{Z}_\bullet)$ is acyclic under our hypothesis. Thus $(\mathcal{Z}_\bullet)$ is a resolution of $H_0(\mathcal{Z}_\bullet) \simeq \mathcal{S}_\mathcal{I}$.

Theorem 3. *Let $I = (f_0, \dots, f_3)$ be a homogeneous ideal of R such that $\text{indeg}(I^{\text{sat}}) = \text{indeg}(I) = d$. Suppose that $\mathcal{P} = \text{Proj}(R/I)$ is locally a complete intersection of dimension zero. Then there exists a complex $(C_\bullet)$ of free B -modules*

$$\begin{array}{ccccccc} 0 & \longrightarrow & B(-3)^l & \longrightarrow & B(-2)^m & \longrightarrow & B(-1)^n \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ & & C_3 & & C_2 & & C_1 \end{array}$$

where $n = \dim_k H_{\mathfrak{m}}^3(Z_1)_{d-1}$, $m = \dim_k H_{\mathfrak{m}}^3(Z_2)_{2d-1}$, $l = \dim_k H_{\mathfrak{m}}^3(Z_3)_{3d-1}$ with homology

$$H_1(C_\bullet) \simeq \oplus_{s \geq 0} H_{\mathfrak{m}}^2(I^s)_{sd-1}, \quad H_2(C_\bullet) \simeq \oplus_{s \geq 0} (I^s)_{sd-1}^{\text{sat}}, \quad H_3(C_\bullet) = 0.$$

Proof. One has the following graded (degree zero) isomorphisms of R -modules (see [BC05, Lemma 1]):

$$H_{\mathfrak{m}}^p(Z_q) \simeq \begin{cases} 0 & \text{for } p = 0, 1 \\ H_{\mathfrak{m}}^0(H_{p-q})^*[3-4d] & \text{for } p = 2 \\ Z_{p-q}^*[3-4d] & \text{for } p = 3 \end{cases} \quad (1)$$

where $-^* := \text{Hom}_{\text{gr}_R}(-, k)$.

We consider the two spectral sequences associated to the double complex $C_q^p = C_{\mathfrak{m}}^p(\mathcal{Z}_q)$, where $C_{\mathfrak{m}}^\bullet(M)$ denotes the Čech complex on M relatively to the ideal $\mathfrak{m}$. Since $(\mathcal{Z}_\bullet)$ is acyclic, one of them abuts at step two with:

$${}_\infty \mathbf{E}_q^p = {}_2 \mathbf{E}_q^p = \begin{cases} H_{\mathfrak{m}}^p(\mathcal{S}_\mathcal{I}), & \text{for } q = 0 \\ 0, & \text{for } q \neq 0. \end{cases} \quad (2)$$

The other one gives at step one:

$${}_1 \mathbf{E}_q^p = H_{\mathfrak{m}}^p(Z_q) = H_{\mathfrak{m}}^p(Z_q)[qd] \otimes_R R[T_0, \dots, T_3](-q) = H_{\mathfrak{m}}^p(Z_q)[qd] \otimes_k B(-q).$$

Notice that $H_{\mathfrak{m}}^p(Z_q) = 0$ for $p = 0, 1$; $Z_3 \simeq R[-4d]$ and $Z_0 = R$. Therefore the first page of the vertical spectral sequence has only two nonzero lines

$$0 \longrightarrow H_{\mathfrak{m}}^2(Z_2)[2d] \otimes_k B(-2) \rightarrow H_{\mathfrak{m}}^2(Z_1)[d] \otimes_k B(-1) \longrightarrow 0$$

$$H_{\mathfrak{m}}^3(Z_3)[3d] \otimes_k B(-3) \rightarrow H_{\mathfrak{m}}^3(Z_2)[2d] \otimes_k B(-2) \rightarrow H_{\mathfrak{m}}^3(Z_1)[d] \otimes_k B(-1) \rightarrow H_{\mathfrak{m}}^3(Z_0) \otimes_k B.$$

In bidegree $(-1, *)$, we have $H_{\mathfrak{m}}^3(Z_0)_{-1} \otimes_k B = H_{\mathfrak{m}}^3(R)_{-1} \otimes_k B = 0$. Therefore, we obtain the complex $(C_{\bullet})$ of free B -modules

$$\begin{array}{ccccccc} 0 & \longrightarrow & B(-3)^l & \longrightarrow & B(-2)^m & \longrightarrow & B(-1)^n \longrightarrow 0. \\ & & \parallel & & \parallel & & \parallel \\ & & C_3 & & C_2 & & C_1 \end{array}$$

Let us compute the homology of $(C_{\bullet})$. The exact sequence

$$0 \longrightarrow Z_1 \longrightarrow R[-d]^4 \longrightarrow I \longrightarrow 0 \quad (3)$$

gives

$$H_{\mathfrak{m}}^2(Z_1) \simeq H_{\mathfrak{m}}^1(I) \simeq H_{\mathfrak{m}}^0(R/I) \simeq I^{\text{sat}}/I.$$

Since $\text{indeg}(I^{\text{sat}}) = \text{indeg}(I) = d$, we have

$$H_{\mathfrak{m}}^2(Z_1)_{d-1} \simeq (I^{\text{sat}}/I)_{d-1} = 0. \quad (4)$$

It follows from (1) that $H_{\mathfrak{m}}^2(Z_2)_{2d-1} \simeq H_{\mathfrak{m}}^0(H_0)_{2-2d}^* \simeq \text{Homgr}_r((I^{\text{sat}}/I)_{2d-2}, k)$ as $H_{\mathfrak{m}}^0(H_0) \simeq I^{\text{sat}}/I$. Since $\text{reg}(I) \leq 2d-2$ (see [BBC14, Lemma 4]), we have $I_{2d-2} = (I^{\text{sat}})_{2d-2}$, thus

$$H_{\mathfrak{m}}^2(Z_2)_{2d-1} = 0. \quad (5)$$

It follows from (4) and (5) that

$$({}_1^v \mathbf{E}_q^p)_{(-1,*)} = 0 \quad \text{unless } p = 3 \text{ and } q \in \{1, 2, 3\}.$$

Therefore, in bidegree $(-1, *)$,

$${}_{\infty}^v \mathbf{E}_q^p = {}_2^v \mathbf{E}_q^p = \begin{cases} H_q(C_{\bullet}), & \text{if } p = 3 \text{ and } q = 1, 2, 3 \\ 0, & \text{else.} \end{cases} \quad (6)$$

By comparing the two spectral sequences (2) and (6), one has:

$$\begin{aligned} H_1(C_{\bullet}) &= H_{\mathfrak{m}}^2(\mathcal{S}_{\mathcal{I}})_{(-1,*)} = \oplus_{s \geq 0} H_{\mathfrak{m}}^2(I^s)_{sd-1}, \\ H_2(C_{\bullet}) &= H_{\mathfrak{m}}^1(\mathcal{S}_{\mathcal{I}})_{(-1,*)} = \oplus_{s \geq 0} H_{\mathfrak{m}}^1(I^s)_{sd-1} = \oplus_{s \geq 0} H_{\mathfrak{m}}^0(R/I^s)_{sd-1} = \oplus_{s \geq 0} ((I^s)^{\text{sat}}/I^s)_{sd-1} \\ &= \oplus_{s \geq 0} (I^s)_{sd-1}^{\text{sat}} \text{ since } I_{sd-1}^s = 0, \\ H_3(C_{\bullet}) &= H_{\mathfrak{m}}^0(\mathcal{S}_{\mathcal{I}})_{(-1,*)} \subseteq (\mathcal{S}_{\mathcal{I}})_{(-1,*)} \subseteq R_{-1} \otimes_k B = 0. \end{aligned}$$

□

We now give an expression of m, n, l in terms of the degree d and $\deg(\mathcal{P})$.

Lemma 4. *Under the assumptions of Theorem 3,*

$$n = \deg(\mathcal{P}) - \frac{1}{2}d(d+1), \quad m = \deg(\mathcal{P}) - d \text{ and } l = \frac{1}{2}d(d-1).$$

As a consequence, $n - m + l = 0$.

Proof. We first compute $n = \dim_k H_{\mathfrak{m}}^3(Z_1)_{d-1}$. From (3), we derive an exact sequence

$$0 \longrightarrow H_{\mathfrak{m}}^2(I) \longrightarrow H_{\mathfrak{m}}^3(Z_1) \longrightarrow H_{\mathfrak{m}}^3(R)[-d]^4 \longrightarrow 0,$$

hence $H_{\mathfrak{m}}^3(Z_1)_{d-1} \simeq H_{\mathfrak{m}}^2(I)_{d-1}$. We get

$$n = \dim_k H_{\mathfrak{m}}^2(I)_{d-1} = \dim_k H_{\mathfrak{m}}^1(R/I)_{d-1}.$$

We now consider an exact sequence

$$0 \longrightarrow R/I^{\text{sat}} \longrightarrow \bigoplus_{\mu \in \mathbb{Z}} H^0(\mathbb{P}_k^2, \mathcal{O}_{\mathcal{P}}(\mu)) \longrightarrow H_{\mathfrak{m}}^1(R/I) \longrightarrow 0,$$

hence

$$\dim_k H_{\mathfrak{m}}^1(R/I)_{d-1} = \dim_k H^0(\mathbb{P}_k^2, \mathcal{O}_{\mathcal{P}}(d-1)) - \dim_k (R/I^{\text{sat}})_{d-1} = \deg(\mathcal{P}) - \dim_k (R/I^{\text{sat}})_{d-1}.$$

Since $\text{indeg}(I^{\text{sat}}) = d$, $\dim_k (R/I^{\text{sat}})_{d-1} = \dim_k R_{d-1} = \binom{d+1}{2} = \frac{1}{2}d(d+1)$. It follows that

$$n = \deg(\mathcal{P}) - \frac{1}{2}d(d+1).$$

We next compute $m = \dim_k H_{\mathfrak{m}}^3(Z_2)_{2d-1}$. It follows from (1) that

$$H_{\mathfrak{m}}^3(Z_2)_{2d-1} \simeq (Z_1)_{2-2d}^* \simeq \text{Homgr}_R((Z_1)_{2d-2}, k).$$

Therefore, by (3) we have

$$m = \dim_k (Z_1)_{2d-2} = \dim_k R_{d-2}^4 - \dim_k I_{2d-2} = 4 \binom{d}{2} - \dim_k I_{2d-2}.$$

Since $\dim_k I_{2d-2} = \dim_k R_{2d-2} - \dim_k (R/I)_{2d-2} = \binom{2d}{2} - \deg(\mathcal{P})$, it follows that

$$m = 4 \binom{d}{2} - \binom{2d}{2} + \deg(\mathcal{P}) = \deg(\mathcal{P}) - d.$$

Finally, we have

$$l = \dim_k H_{\mathfrak{m}}^3(Z_3)_{3d-1} = \dim_k H_{\mathfrak{m}}^3(R)_{-d-1} = \binom{d}{2} = \frac{1}{2}d(d-1).$$

□

Now we will give a bound for the degree of the projective variety $\mathcal{P} \subset \mathbb{P}_k^2$ in terms of the degree d . Let us introduce some classical notations.

For a finitely generated graded R -module M , we denote by HF_M, HP_M its Hilbert function and its Hilbert polynomial, respectively. By definition, for every μ , $HF_M(\mu) = \dim_k (M_{\mu})$ and by the Grothendieck-Serre formula [BH93, Theorem 4.3.5]

$$HP_M(\mu) = HF_M(\mu) - \sum_i (-1)^i HF_{H_{\mathfrak{m}}^i(M)}(\mu).$$

Lemma 5. *Under the assumptions of Theorem 3,*

$$\frac{1}{2}d(d+1) \leq \deg(\mathcal{P}) \leq d^2 - 2d + 3.$$

Proof. The Hilbert function of R/I^{sat} is weakly increasing, hence for any μ

$$\dim_k (R/I^{\text{sat}})_{\mu} \leq \deg(\mathcal{P}).$$

Since $\text{indeg}(I^{\text{sat}}) = d$, $\deg(\mathcal{P}) \geq \dim_k (R/I^{\text{sat}})_{d-1} = \dim_k R_{d-1} = \frac{1}{2}d(d+1)$.

We can assume that k is infinite. Since $\text{codim}(I) = 2$, there are homogeneous polynomials $g_1, g_2 \in I$ of degree d which form a regular sequence. Therefore $\mathfrak{b} := (g_1, g_2)$ is a complete intersection ideal. Setting $J := \mathfrak{b} :_R I$, J is saturated and $I^{\text{sat}} = \mathfrak{b} :_R J$ since R is Gorenstein. We set $\mathcal{Q} = \text{Proj}(R/J)$. By using liaison theory, we obtain $\deg(\mathcal{P}) + \deg(\mathcal{Q}) = \deg(\mathfrak{b})$, hence $\deg(\mathcal{P}) = d^2 - \deg(\mathcal{Q})$.

Local duality and the liaison sequence show that

$$H_{\mathfrak{m}}^1(R/J) \simeq (I^{\text{sat}}/\mathfrak{b})^*[3-2d].$$

We thus get

$$H_{\mathfrak{m}}^1(R/J)_{\mu} \simeq \text{Homgr}_R((I^{\text{sat}}/\mathfrak{b})_{2d-3-\mu}, k).$$

Since $\text{indeg}(I^{\text{sat}}) = \text{indeg}(\mathfrak{b}) = d$, we obtain

$$H_{\mathfrak{m}}^1(R/J)_{\mu} = \begin{cases} 0, & \text{if } \mu > d-3 \\ \text{Homgr}_R((I^{\text{sat}}/\mathfrak{b})_d, k) \neq 0, & \text{if } \mu = d-3. \end{cases}$$

Moreover, we have $H_{\mathfrak{m}}^0(R/J) = J^{\text{sat}}/J = 0$. We therefore conclude that

$$\text{reg}(R/J) = d-2,$$

and hence $\dim_k(R/J)_{d-2} = \deg(\mathcal{Q})$.

On the other hand, the Hilbert function $HF_{R/J}(\mu)$ of R/J strictly increases from 1 in degree 0 until $\deg(\mathcal{Q})$ in degree $d-2$. Therefore one has $\deg(\mathcal{Q}) \geq d-1$. By the Grothendieck-Serre formula, for any μ

$$HF_{R/J}(\mu) = \deg(\mathcal{Q}) - \dim_k H_{\mathfrak{m}}^1(R/J)_{\mu}.$$

Since $\dim_k H_{\mathfrak{m}}^1(R/J)_{d-3} = \dim_k(I^{\text{sat}}/\mathfrak{b})_d \geq 2$, we deduce $\deg(\mathcal{Q}) \geq d$.

The proof will be completed by showing that $\deg(\mathcal{Q}) \geq 2d-3$. Set $c_i = \dim_k(R/J)_i$ for all $i \geq 0$. The sequence $\underline{c} = \{c_i : i \geq 0\}$ satisfies the following properties:

- (i) $c_0 = 1$ and $c_1 = 3$. Indeed, if $c_1 < 3$, then $J_1 \neq 0$, there is therefore a linear form $\ell \in J_1$. We assume that $J = (\ell, w_1, w_2, \dots, w_s)$. Since $\text{reg}(J) = d-1$, we have $\deg(w_i) \leq d-1$ for all $i = 1, \dots, s$. Fix $i \in \{1, \dots, s\}$ such that $\gcd(\ell, w_i) = 1$. Since $\mathcal{Q} \subseteq V(\ell, w_i)$, hence $\deg(\mathcal{Q}) \leq d-1$, a contradiction.
- (ii) $c_i = \deg(\mathcal{Q})$ for any $i \geq d-2$ and $c_{d-2} - c_{d-3} = \dim_k H_{\mathfrak{m}}^1(R/J)_{d-3} \geq 2$.

Since J is saturated and k is infinite, there is a linear form ℓ in R such that $J : \ell \simeq J(-1)$ as graded R -modules. It follows that ℓ induces an injection $\times \ell : (R/J)[-1] \rightarrow R/J$. Therefore the Hilbert function of $R/(J+\ell)$ is $\underline{b} = \{b_i : i \geq 0\}$ with $b_0 = 1$ and $b_i = c_i - c_{i-1}$ for every $i \geq 1$. Since $b_{d-2} \geq 2$, as a consequence of Macaulay's Theorem $b_i \geq 2$, for any $i = 1, \dots, d-2$.

Indeed, suppose that there exists an integer $i \geq 1$ such that $b_i = 1$, and let

$$\nu = \sup_{i \geq 1} \{i \mid b_i = 1\}.$$

One has $2 \leq \nu \leq d-3$ and since $R/(J+\ell)$ is a standard graded k -algebra, Macaulay's Theorem (see [Sta78, Theorem 2.2]) shows that

$$2 \leq b_{\nu+1} \leq b_{\nu}^{<\nu>} = 1^{<\nu>} = 1.$$

This shows that $c_{i+1} \geq c_i + 2$ for all $i = 0, \dots, d-3$, hence

$$\deg(\mathcal{Q}) \geq 2(d-2) + 1 = 2d-3.$$

□

Proposition 6. *Under the assumptions of Theorem 3, the following holds for any integer s :*

$$\dim_k H_{\mathfrak{m}}^2(I^s)_{sd-1} = \frac{1}{2}s(s+1)\deg(\mathcal{P}) - \frac{1}{2}sd(sd+1) + \dim_k(I^s)_{sd-1}^{\text{sat}}.$$

Proof. By Theorem 3, for any integer s , one has

$$\begin{aligned}
\dim_k H_{\mathfrak{m}}^2(I^s)_{sd-1} &= n \cdot \dim_k B_{s-1} - m \cdot \dim_k B_{s-2} + l \cdot \dim_k B_{s-3} + \dim_k (I^s)_{sd-1}^{\text{sat}} \\
&= n \binom{s+2}{3} - (n+l) \binom{s+1}{3} + l \binom{s}{3} + \dim_k (I^s)_{sd-1}^{\text{sat}} \\
&= n \binom{s+1}{2} - l \binom{s}{2} + \dim_k (I^s)_{sd-1}^{\text{sat}} \\
&= \left(\deg(\mathcal{P}) - \frac{1}{2}d(d+1) \right) \binom{s+1}{2} - \frac{1}{2}d(d-1) \binom{s}{2} + \dim_k (I^s)_{sd-1}^{\text{sat}} \\
&= \frac{1}{2}s(s+1) \deg(\mathcal{P}) - \frac{1}{2}sd(sd+1) + \dim_k (I^s)_{sd-1}^{\text{sat}}.
\end{aligned}$$

□

Theorem 7. *Let I be a homogeneous ideal of R such that $\text{indeg}(I^{\text{sat}}) = \text{indeg}(I) = 3$. If $\mathcal{P} = \text{Proj}(R/I)$ is locally a complete intersection of dimension zero, then*

$$\sum_{\mathfrak{p} \in \mathcal{Y}} \deg(h_{\mathfrak{p}}) \leq 4.$$

Proof. We will prove first that $I = I^{\text{sat}}$. In this case, $\deg(\mathcal{P}) = 6$ by Lemma 5. Moreover, with the notations in the proof of Lemma 5, we have

$$H_{\mathfrak{m}}^1(R/I^{\text{sat}}) \simeq (J/\mathfrak{b})^*[-3]$$

and $\text{reg}(R/J) = 1$. Hence $\dim_k(R/J)_{\mu} = \deg(\mathcal{Q}) = 3$ for all $\mu \geq 1$. It follows that

$$\dim_k H_{\mathfrak{m}}^1(R/I^{\text{sat}})_{\mu} = 0, \forall \mu \geq 2, \dim_k H_{\mathfrak{m}}^1(R/I^{\text{sat}})_1 = 3.$$

Therefore $\text{reg}(R/I^{\text{sat}}) = 2$, hence I^{sat} is generated in degree at most 3. Since $\text{indeg}(I^{\text{sat}}) = 3$, it is immediate that I^{sat} is exactly generated in degree 3. Furthermore,

$$\dim_k(I^{\text{sat}})_3 = \dim_k R_3 - \dim_k(R/I^{\text{sat}})_3 = 10 - 6 = 4,$$

which implies $I^{\text{sat}} = I$.

By the Hilbert-Burch Theorem, I admits a free resolution of the form

$$0 \longrightarrow R(-4)^3 \xrightarrow{M} R(-3)^4 \longrightarrow R \longrightarrow R/I \longrightarrow 0.$$

By changing the coordinates, there is no loss of generality in assuming the matrix M be of the form

$$M = \begin{pmatrix} X_1 & 0 & 0 \\ a_1 X_2 & X_2 & 0 \\ b_1 X_3 & b_2 X_3 & X_3 \\ c_1 L & c_2 L & c_3 L \end{pmatrix}$$

where $L = \alpha_1 X_1 + \alpha_2 X_2 + \alpha_3 X_3$. One has $c_3 \neq 0$ since $f_i \neq 0$ for all $i = 0, 1, 2, 3$. Since I is locally a complete intersection, $V(I_2(M)) = \emptyset$, where $I_2(M)$ is the ideal of (2×2) -minors of M . It follows that $\alpha_1 \alpha_2 \alpha_3 \neq 0$.

Let now a fiber over a closed point $\mathfrak{p}$ of coordinates $(\lambda_1 : \lambda_2 : \lambda_3 : \lambda_4) \in \mathbb{P}_k^3$ be of dimension 1 and its unmixed component be defined by $h_{\mathfrak{p}} \in R$. It follows from [BBC14, Lemma 10] that

$$h_{\mathfrak{p}} = \gcd(\ell_1, \ell_2, \ell_3),$$

where

$$\begin{aligned}\ell_1 &= (\lambda_1 + c_1\alpha_1\lambda_4)X_1 + (a_1\lambda_2 + c_1\alpha_2\lambda_4)X_2 + (b_1\lambda_3 + c_1\alpha_3\lambda_4)X_3, \\ \ell_2 &= c_2\alpha_1\lambda_4X_1 + (\lambda_2 + c_2\alpha_2\lambda_4)X_2 + (b_2\lambda_3 + c_2\alpha_3\lambda_4)X_3, \\ \ell_3 &= c_3\alpha_1\lambda_4X_1 + c_3\alpha_2\lambda_4X_2 + (\lambda_3 + c_3\alpha_3\lambda_4)X_3.\end{aligned}$$

- If $\lambda_4 = 0$, then there are the following cases:

Case 1: $\lambda_3 \neq 0$. In this case we have $\lambda_1 = \lambda_2 = 0$. Therefore $\mathbf{p} = (0 : 0 : 1 : 0)$ and $h_{\mathbf{p}} = X_3$.

Case 2: $\lambda_3 = 0$ and $\lambda_2 \neq 0$. It follows that $\lambda_1 = 0$. Thus $\mathbf{p} = (0 : 1 : 0 : 0)$ and $h_{\mathbf{p}} = X_2$.

Case 3: $\lambda_3 = 0$ and $\lambda_2 = 0$. We must have $\lambda_1 \neq 0$. Hence $\mathbf{p} = (1 : 0 : 0 : 0)$ and $h_{\mathbf{p}} = X_1$.

- If $\lambda_4 = 1$, then from three polynomials ℓ_1, ℓ_2, ℓ_3 have a common divisor, we have

$$\begin{aligned}\frac{\lambda_1 + c_1\alpha_1}{c_3\alpha_1} &= \frac{a_1\lambda_2 + c_1\alpha_2}{c_3\alpha_2} = \frac{b_1\lambda_3 + c_1\alpha_3}{\lambda_3 + c_3\alpha_3}, \\ \frac{c_2}{c_3} &= \frac{\lambda_2 + c_2\alpha_2}{c_3\alpha_2} = \frac{b_2\lambda_3 + c_2\alpha_3}{\lambda_3 + c_3\alpha_3}.\end{aligned}$$

These equalities imply that $\lambda_1 = \lambda_2 = 0$. Furthermore, we also have $\lambda_3 = 0$. Indeed, if $\lambda_3 \neq 0$, then $c_1 = b_1c_3$ and $c_2 = b_2c_3$, contrary to $f_i \neq 0$ for all $i = 0, \dots, 3$. Therefore $\mathbf{p} = (0 : 0 : 0 : 1)$ and $h_{\mathbf{p}} = L$. $\square$

Example 8. Consider the parameterization given by

$$\begin{aligned}f_0 &= X_2X_3(X_1 + X_2 + X_3), \\ f_1 &= X_1X_3(X_1 + X_2 + X_3), \\ f_2 &= X_1X_2(X_1 + X_2 + X_3), \\ f_3 &= X_1X_2X_3.\end{aligned}$$

Using `Macaulay2` [Mac2], we see that $\mathcal{P}$ is locally a complete intersection of degree 6 and that I is a saturated ideal with free resolution

$$0 \longrightarrow R(-4)^3 \xrightarrow{M} R(-3)^4 \longrightarrow R \longrightarrow R/I \longrightarrow 0,$$

where the matrix M is given by

$$\begin{pmatrix} -X_3 & 0 & 0 \\ X_2 & -X_2 & 0 \\ 0 & X_1 + X_2 + X_3 & -X_2 - X_3 \\ 0 & 0 & X_1 \end{pmatrix}.$$

Thus we obtain $\mathcal{Y} = \{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4\}$ with

$$\begin{aligned}\mathbf{p}_1 &= (1 : 0 : 0 : 0), & h_{\mathbf{p}_1} &= X_3, & \mathbf{p}_2 &= (0 : 1 : 0 : 0), & h_{\mathbf{p}_2} &= X_2, \\ \mathbf{p}_3 &= (0 : 0 : 0 : 1), & h_{\mathbf{p}_3} &= X_1, & \mathbf{p}_4 &= (0 : 0 : 1 : -1), & h_{\mathbf{p}_4} &= X_1 + X_2 + X_3.\end{aligned}$$

Consequently, we have

$$\sum_{\mathbf{p} \in \mathcal{Y}} \deg(h_{\mathbf{p}}) = 4.$$

The following example shows that the hypothesis I to be locally a complete intersection outside $V(\mathfrak{m})$ is necessary.

Example 9. Consider the parameterization given by

$$\begin{aligned} f_0 &= (X_1 + X_2 + X_3)(X_1 + 2X_2 + 2X_3)(X_2 + X_3), \\ f_1 &= X_1(X_1 - X_2 - X_3)(X_2 + X_3), \\ f_2 &= X_1(X_1 + X_2 + X_3)(X_1 + 2X_2 + 2X_3), \\ f_3 &= X_1(X_1 + X_2 + X_3)(X_1 - X_2 - X_3). \end{aligned}$$

Using `Macaulay2` [Mac2], we deduce that $\mathcal{P}$ is not locally a complete intersection of degree 6 and that I is a saturated ideal with free resolution

$$0 \longrightarrow R(-4)^3 \xrightarrow{M} R(-3)^4 \longrightarrow R \longrightarrow R/I \longrightarrow 0,$$

where the matrix M is given by

$$M = \begin{pmatrix} -X_2 - X_3 & 0 & 0 \\ X_1 - X_2 - X_3 & -X_2 - X_3 & 0 \\ 0 & X_1 + X_2 + X_3 & -X_2 - X_3 \\ 0 & 0 & X_1 \end{pmatrix}.$$

Hence $\mathcal{Y} = \{\mathfrak{p}_1, \mathfrak{p}_2, \mathfrak{p}_3, \mathfrak{p}_4, \mathfrak{p}_5\}$ with

$$\begin{aligned} \mathfrak{p}_1 &= (1 : 0 : 0 : 0), & h_{\mathfrak{p}_1} &= X_2 + X_3, & \mathfrak{p}_2 &= (0 : 0 : 0 : 1), & h_{\mathfrak{p}_2} &= X_1, \\ \mathfrak{p}_3 &= (0 : 0 : 1 : -1), & h_{\mathfrak{p}_3} &= X_1 + X_2 + X_3, & \mathfrak{p}_4 &= (0 : 2 : 1 : 1), & h_{\mathfrak{p}_4} &= X_1 - X_2 - X_3, \\ \mathfrak{p}_5 &= (6 : -2 : 2 : -1), & h_{\mathfrak{p}_5} &= X_1 + 2X_2 + 2X_3. \end{aligned}$$

It follows that

$$\sum_{\mathfrak{p} \in \mathcal{Y}} \deg(h_{\mathfrak{p}}) = 5 > 4.$$

Theorem 10. Let I be a homogeneous ideal of R such that $\text{indeg}(I^{\text{sat}}) = \text{indeg}(I) = d \geq 4$. If $\mathcal{P} = \text{Proj}(R/I)$ is locally a complete intersection of dimension zero, then

$$\sum_{\mathfrak{p} \in \mathcal{Y}} \deg(h_{\mathfrak{p}}) \leq \left\lfloor \frac{d}{2} \right\rfloor d - 1$$

where $\lfloor x \rfloor := \max\{n \in \mathbb{Z} \mid n \leq x\}$.

Proof. Set $\nu = \lfloor \frac{d}{2} \rfloor$. By Proposition 1, it suffices to show that $(I^\nu)_{\nu d - 1}^{\text{sat}} \neq 0$. Suppose on the contrary that $(I^\nu)_{\nu d - 1}^{\text{sat}} = 0$. By Proposition 6, one has

$$\dim_k H_{\mathfrak{m}}^2(I^\nu)_{\nu d - 1} = \frac{1}{2}\nu(\nu + 1)\deg(\mathcal{P}) - \frac{1}{2}\nu d(\nu d + 1).$$

We now show that this is not possible since $\deg(\mathcal{P}) \leq d^2 - 2d + 3$ by Lemma 5. Indeed, consider two cases:

(i) If $d = 2s$ with $s \geq 2$, then $\nu = \lfloor \frac{d}{2} \rfloor = s$. Therefore

$$\begin{aligned} \dim_k H_{\mathfrak{m}}^2(I^s)_{sd - 1} &= \frac{1}{2}s((s + 1)\deg(\mathcal{P}) - 2s(2s^2 + 1)) \\ &\leq \frac{1}{2}s((s + 1)(4s^2 - 4s + 3) - 2s(2s^2 + 1)) \\ &\leq -\frac{3}{2}s(s - 1) < 0, \end{aligned}$$

which is impossible.

- (ii) Similarly, if $d = 2s+1$ with $s \geq 2$, then $\nu = \lfloor \frac{d}{2} \rfloor = s$. Thus $\dim_k H_m^2(I^s)_{sd-1} \leq -\frac{1}{2}s(s-1) < 0$, which is impossible.

□

Example 11. Let $d \geq 4$ be an integer. Consider the parameterization given by

$$\begin{aligned} f_0 &= X_1^{d-3} X_2 (X_1^2 - X_2^2), \\ f_1 &= X_1^{d-3} X_3 (X_1^2 - X_2^2), \\ f_2 &= X_1^{d-3} X_3 (X_2^2 - X_3^2), \\ f_3 &= X_2^{d-3} X_3 (X_2^2 - X_3^2). \end{aligned}$$

We see that $\mathcal{P}$ is locally a complete intersection of degree $d^2 - 3d + 7$. Moreover, I is a saturated ideal with free resolution

$$0 \longrightarrow R(-d-1) \oplus R(-d-2) \oplus R(-2d+3) \xrightarrow{M} R(-d)^4 \longrightarrow R \longrightarrow R/I \longrightarrow 0,$$

where the matrix M is given by

$$\begin{pmatrix} -X_3 & 0 & 0 \\ X_2 & X_2^2 - X_3^2 & 0 \\ 0 & X_1^2 - X_2^2 & X_2^{d-3} \\ 0 & 0 & X_1^{d-3} \end{pmatrix}.$$

We consider the following cases.

Case 1: if $d-3$ is odd, we obtain $\mathcal{Y} = \{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4, \mathbf{p}_5, \mathbf{p}_6\}$ with

$$\begin{aligned} \mathbf{p}_1 &= (1 : 0 : 0 : 0), & h_{\mathbf{p}_1} &= X_3, & \mathbf{p}_2 &= (0 : 0 : 0 : 1), & h_{\mathbf{p}_2} &= X_1^{d-3}, \\ \mathbf{p}_3 &= (1 : 1 : 0 : 0), & h_{\mathbf{p}_3} &= X_2 - X_3, & \mathbf{p}_4 &= (1 : -1 : 0 : 0), & h_{\mathbf{p}_4} &= X_2 + X_3, \\ \mathbf{p}_5 &= (0 : 0 : 1 : 1), & h_{\mathbf{p}_5} &= X_1 + X_2, & \mathbf{p}_6 &= (0 : 0 : 1 : -1), & h_{\mathbf{p}_6} &= X_1 - X_2. \end{aligned}$$

Case 2: if $d-3$ is even, we have $\mathcal{Y} = \{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4, \mathbf{p}_5\}$ with

$$\begin{aligned} \mathbf{p}_1 &= (1 : 0 : 0 : 0), & h_{\mathbf{p}_1} &= X_3, & \mathbf{p}_2 &= (0 : 0 : 0 : 1), & h_{\mathbf{p}_2} &= X_1^{d-3}, \\ \mathbf{p}_3 &= (1 : 1 : 0 : 0), & h_{\mathbf{p}_3} &= X_2 - X_3, & \mathbf{p}_4 &= (1 : -1 : 0 : 0), & h_{\mathbf{p}_4} &= X_2 + X_3, \\ \mathbf{p}_5 &= (0 : 0 : 1 : -1), & h_{\mathbf{p}_5} &= X_1^2 - X_2^2. \end{aligned}$$

Therefore

$$\sum_{\mathbf{p} \in \mathcal{Y}} \deg(h_{\mathbf{p}}) = d + 2.$$

Question 12. Is $\sum_{\mathbf{p} \in \mathcal{Y}} \deg(h_{\mathbf{p}})$ bounded linearly in the degree d ?

ACKNOWLEDGMENTS

This work was done as a part of the authors Ph.D. thesis at Pierre and Marie Curie University (UPMC), France. He would like to thank his advisor, Professor Marc Chardin, for suggesting the problem, for his enormous help, for being so supportive and his careful reading of the manuscript.

REFERENCES

- [BBC14] Nicolás Botbol and Laurent Busé and Marc Chardin. *Fitting ideals and multiple points of surface parameterizations*. Journal of Algebra, **420**:486–508, 2014.
- [BC05] Laurent Busé and Marc Chardin. *Implicitizing rational hypersurfaces using approximation complexes*. Journal of Symbolic Computation, **40**(4-5):1150–1168, 2005.
- [BCD03] Laurent Busé, David Cox, and Carlos D’Andrea. *Implicitization of surfaces in $\mathbb{P}^3$ in the presence of base points*. J. Algebra Appl., **2**(2):189–214, 2003.
- [BCJ09] Laurent Busé and Marc Chardin and Jean-Pierre Jouanolou. *Torsion of the symmetric algebra and implicitization*. Pro. Amer. Math. Soc., **137**(6):189–214, 2009.
- [BH93] Winfried Bruns and Jürgen Herzog. *Cohen-Macaulay rings*. Volume 39 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1993.
- [BJ03] Laurent Busé and Jean-Pierre Jouanolou. *On the closed image of a rational map and the implicitization problem*. J. Algebra, **265**(1):312–357, 2003.
- [Bus01] Laurent Busé. *Etude du resultant sur une variété algébrique*. Ph.D. Thesis, Université de Nice Sophia-Antipolis, 2001.
- [Bus14] Laurent Busé. *Implicit matrix representations of rational Bézier curves and surfaces*. Computer-Aided Design, Elsevier, 2014, 2013 SIAM Conference on Geometric and Physical Modeling, J. Algebra, **46**(hal-00847802):14–24, 2014.
- [Cha06] Marc Chardin. *Implicitization using approximation complexes*. Algebraic geometry and geometric modeling. Part of the series Mathematics and Visualization, 23–35, 2006.
- [Cox01] David A Cox. *Equations of parametric curves and surfaces via syzygies*. Symbolic Computation: Solving Equations in Algebra, Geometry, and Engineering. Contemporary Mathematics, Amer. Math. Soc., **286**:1–20, 2001.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Graduate Texts in Mathematics 52. Springer-Verlag, New York-Heidelberg, 1977.
- [Hof89] Christoph M. Hoffmann. *Geometric and Solid Modeling: an Introduction*. Morgan Kaufmann Publishers, Inc., 1989.
- [Jou96] Jean-Pierre Jouanolou. *Resultant anisotrope: Compléments et applications*. Electron. J. Combin. 3, Research Paper 2, 1996.
- [Mac2] Daniel R. Grayson and Michael E. Stillman. Macaulay 2, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [MC92] Dinesh Manocha and John F. Canny. *Algorithm for implicitizing rational parametric surfaces*. Comput. Aided Geom. Design, **9**(1):25–50, 1992.
- [SC95] Tom Sederberg and Falai Chen. *Implicitization using moving curves and surfaces*. Computer Graphics Annual Conference Series, (303):301–308, 1995.
- [Sta78] Richard P. Stanley. *Hilbert functions of graded algebras*. Advances in Mathematics, (28):57–83, 1978.
- [ZS60] Oscar Zariski and Pierre Samuel. *Commutative Algebra with the cooperation of I.S. Cohen*. Volume II of Graduate Texts in Mathematics, Springer-Verlag, New York-Heidelberg, 1960.

HUE UNIVERSITY’S COLLEGE OF EDUCATION, 34 LE LOI ST., HUE CITY, VIETNAM

CURRENT ADDRESS: INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UNIVERSITÉ PIERRE ET MARIE CURIE, BOITE THÉSARD, 4 PLACE JUSSIEU, F-75252 PARIS CEDEX 05, FRANCE

E-mail address: quang-hoa.tran@imj-prg.fr